

Docket No. 2251/76935/JPW/GJG/REB

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Yingru Wu, et al. Examiner: Stuart F. Baum
U.S. Serial No.: 10/594,785 Group Art: 1638
Filed : September 17, 2007
For : GENES INVOLVED IN PLANT FIBRE DEVELOPMENT

30 Rockefeller Plaza
New York, New York 10112

BY EFS

Commissioner for Patents
Alexandria, VA 22313-1450

SIR:

DECLARATION UNDER 37 C.F.R. §1.132 OF DANNY J. LLEWELLYN

I, Danny J. Llewellyn, hereby declare that:

1. I am named as an inventor on the above-identified patent application.
2. I am presently a Sub-Program Leader of Genomics and Plant Development at the Commonwealth Scientific and Industrial Research Organisation, Black Mountain Laboratories, Clunies Ross St, Black Mountain ACT 2601, Australia. A copy of my curriculum vitae is attached hereto as **Exhibit A**.
3. I have reviewed and am familiar with the subject application, including pending claims 117-142, which are attached here to as **Exhibit B**.

Applicant: Yingru Wu, et al.
Serial No.: 10/594,785
Filed: September 17, 2007
Exhibit 2

Applicants: Yingru Wu, et al.
Serial No.: 10/594,785
Filed : September 17, 2007
Page : 2 of 8

4. The subject application discloses, e.g. in Examples 5, 6, and 14, that *GhMYB25* and *GhMYB25-like* encode MYB proteins that are 69% identical to each other on the amino acid level, both have similar gene expression profiles in wild-type plants, and both have reduced expression in a cotton mutant with reduced fibre production.
5. In Example 12 of the subject application, the function of *GhMYB25* is analyzed and also represents the expected function of *GhMYB25-like*.
6. Example 12 of the subject application describes experiments that show transgenic tobacco plants over-expressing a *GhMYB25* transgene have increased trichome production on their leaves.
7. Example 12 of the subject application also shows the amount of increased trichome production on the leaves of transgenic tobacco plants correlates with the amount of over-expression of a *GhMYB25* transgene in the transgenic tobacco plants.
8. Increased trichome production on the leaves of transgenic tobacco plants over-expressing a *GhMYB25* transgene indicates that such transgene would cause increased fibre initiation and/or elongation if over-expressed in a fibre producing plant.
9. Transgenic tobacco plants over-expressing a *GhMYB25* transgene having increased trichome production on their

Applicants: Yingru Wu, et al.
Serial No.: 10/594,785
Filed : September 17, 2007
Page : 3 of 8

leaves is predictive that a *GhMYB25*-like transgene would cause increased fibre initiation and/or elongation if over-expressed in fibre producing plants.

10. Example 13 of the subject application describes how a transgenic fibre producing plant over-expressing a *GhMYB25*-like transgene may be created.
11. We have confirmed through experiments that the predictions within the subject application, in particular Example 13, were correct. The data discussed below was obtained by me or individuals working under my guidance and supervision.
12. The experiments discussed below in paragraphs 13-20, were performed to determine whether over-expressing *GhMYB25*-like in a transgenic fibre producing plant results in a transgenic plant having increased fibre initiation and/or elongation compared to wild-type plants.
13. The coding region of the D-genome homologue of *GhMYB25*-like driven by either a seed-specific (ovule epidermis) promoter, *FBP7* from petunia, or a constitutive viral promoter, *Stunt7*, were successfully introduced into cotton cultivar Coker 315 by *A. tumefaciens*-mediated transformation. The D-genome homologue of *GhMYB25*-like encodes a polypeptide also encoded by a nucleotide sequence which is identical to the nucleotide sequence set forth in SEQ ID NO: 38 and described in the subject application.

Applicants: Yingru Wu, et al.
Serial No.: 10/594,785
Filed : September 17, 2007
Page : 4 of 8

14. A total of 11 and 36 independent T_0 lines were produced for the *FBP7-GhMYB25-like* and the *S7-GhMYB25-like* constructs, respectively and T_1 seed harvested from 10 independent *FBP7-GhMYB25-like* and 2 independent *S7-GhMYB25-like* transgenic lines.
15. Vegetative growth of the T_1 plants appeared to be the same as wild-type plants.
16. Initial scanning electron microscope (SEM) analysis was performed on T_2 ovules (ovules produced on T_1 plants) from those transgenic lines as available. The number of fibre initiations was counted from SEM images in the middle of 0 dpa ovules and approximately 12 ovules were imaged per line, from multiple plants from each line.
17. The ovules from *FBP7-GhMYB25-like* plants showed an average increase in the number of fibre initiations between 13% and 28% in five independent lines compared to wild-type controls, with some individual ovules having 30-50% more fibre initiations.
18. Of the two *S7-GhMYB25-like* transgenic lines analyzed, one showed an average increase of 13% and the second showed a 22% increase in the number of fibre initiations compared to wild-type controls, again with a number of ovules having 60% more fibre initiations.

Applicants: Yingru Wu, et al.
Serial No.: 10/594,785
Filed : September 17, 2007
Page : 5 of 8

19. These experiments demonstrated that the increased production by a fibre producing plant of a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38 of the subject application results in increased fibre initiation and/or elongation in the fibre producing plant.
20. These experiments also demonstrate the creation of a transgenic fibre producing plant with increased fibre initiation and/or elongation by genetically manipulating the plant such that the production of a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38 of the subject application is increased when compared to a wild-type fibre producing plant.
21. Example 5 and Table 4 of the subject specification show that expression of the *GhMYB25* and *GhMYB25-like* genes is reduced in a mutant fibre producing plant with reduced fibre initiation and/or elongation in the plant.
22. A mutant fibre producing plant with reduced expression of the *GhMYB25-like* gene having reduced fibre initiation and/or elongation is predictive that reduction of the gene in a wild-type plant would cause reduced fibre initiation and/or elongation in that plant.

Applicants: Yingru Wu, et al.
Serial No.: 10/594,785
Filed : September 17, 2007
Page : 6 of 8

23. Example 13 of the subject application describes how a transgenic fibre producing plant with reduced expression of the *GhMYB25*-like gene may be created.
24. We have confirmed that reduced expression of *GhMYB25*-like results in reduced fibre initiation and/or elongation in fibre producing plants. The data discussed below was obtained by me or individuals working under my guidance and supervision.
25. The experiments discussed below in paragraphs 26-32, were performed to determine whether reducing the expression of *GhMYB25*-like in a transgenic fibre producing plant results in a transgenic plant having reduced fibre initiation and/or elongation compared to wild-type plants.
26. A constitutive 35S promoter-driven RNAi construct targeted against both the A and D-genome homologues of *GhMYB25*-like was introduced into cotton cultivar Coker 315 by *A. tumefaciens*-mediated transformation, and seven independent transgenic lines were produced.
27. Detailed phenotypic analysis was performed on four independent *GhMYB25*-like silenced (25Li) lines.
28. For each line, T₃ plants homozygous for the presence of the transgene or wild-type segregants of the same generation were analyzed.

Applicants: Yingru Wu, et al.
Serial No.: 10/594,785
Filed : September 17, 2007
Page : 7 of 8

29. The transcript levels for *GhMYB25-like* in these plants were significantly reduced in the ovules and seeds of all the silenced plants compared to their corresponding wild-type controls.
30. The seeds from the T₃ homozygous 25Li plants were essentially fibreless, except for a few fibres at the chalazal end, when compared to the normal fibred seeds from the wild-type plants. Fuzz fibres were also completely absent from all 25Li plants. The *GhMYB25-like* silenced plants were indistinguishable from their wild-type controls in respect to observable morphological features such as leaf shape, height, flowering time and flower structure.
31. These experiments demonstrated that the reduced production by a fibre producing plant of a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38 of the subject application results in reduced fibre initiation and/or elongation in the fibre producing plant.
32. These experiments also demonstrate the creation of a transgenic fibre producing plant with reduced fibre initiation and/or elongation by genetically manipulating the plant such that the production of a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38 of the

Applicants: Yingru Wu, et al.
Serial No.: 10/594,785
Filed : September 17, 2007
Page : 8 of 8

subject application is reduced when compared to a wild-type fibre producing plant.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made herein on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the subject application or any patent issuing thereon.

Dated: 31/1/11



Danny J. Blewellyn

Curriculum Vitae

Danny James Llewellyn, PhD, FTSE

Professional Address: CSIRO Plant Industry
P.O. Box 1600
Canberra , ACT 2601

Nationality: Australian

Date and place of birth: 21/7/1954 Newcastle, NSW, Australia

Current Position: Group and Project Leader in Cotton Biotechnology, CSIRO
Plant Industry, Canberra

Academic record/awards:

1975	BSc (Hons) Australian National University. Majoring in Biochemistry
1980	PhD Australian National University. Thesis on multifunctional enzymes
1994	Australian Cotton Grower's Research Association Researcher of the Year
1995	CSIRO Plant Industry Chief's Award (Excellence in Science Delivery)
2000	Elected a Fellow of the Australian Academy of Technological Sciences and Engineering (ATSE) for contributions to Australian Agricultural Biotechnology
2001	Centenary Medal for services to Agricultural Biotechnology
2003	CSIRO Plant Industry Chief's Award (Excellence in Research Adoption)
2003	CSIRO Chairman's Medal
2006	ATSE Clunies Ross Award
2009	Weed Science Society of America Award of Excellence

Employment record:

1995 to present:	CSIRO Plant Industry (cotton biotechnology group and project leader)
1993 to 1999	Sub-program Leader Cotton CRC
1985 to 1995	Research Scientist, CSIRO Plant Industry Canberra
1982 to 1985:	Rural Credits Research Fellow, CSIRO Plant Industry, Canberra
1980 to 1982:	CSIRO Postdoctoral Fellow, Max-Planck Institute for Plant Breeding, Cologne, Germany
1980 to 1980:	Research Assistant, Dept Biochemistry, ANU, Canberra – while completing PhD thesis

Professional societies

Australian Society for Biochemistry and Molecular Biology
Australian Academy of Technological Sciences and Engineering

Refereed Publications:

Llewellyn, D.J., and Smith, G.D. (1978) An evaluation of active enzyme centrifugation as a zonal and boundary technique by the analysis of simulated data. *Arch Biochem Biophys.* 190:483-94.

Applicant: Yingru Wu, et al.
Serial No.:10/594,785
Filed: September 17, 2007
Exhibit A

- Llewellyn, D.J. and Smith, G.D. (1979) Study of chorismate mutase--prephenate dehydrogenase in crude cell extracts of *Escherichia coli*. *Biochemistry* **18**:4704-14.
- Llewellyn, D.J., Daday, A. and Smith, G.D. (1980) Evidence for an artificially evolved bifunctional 3-deoxy-D-arabinoheptulosonate-7-phosphate synthase-chorismate mutase in *Bacillus subtilis*. *J Biol Chem.* **255**:2077-84.
- Llewellyn, D.J. and Smith, G.D. (1981) The relationship between molecular weight and quaternary structure of globular proteins. *Arch Biochem Biophys.* **207**:63-8.
- Dennis, E.S., Gerlach, W.L., Pryor, A.J., Bennetzen, J.L., Inglis, A., Llewellyn, D., Sachs, M.M., Ferl, R.J. and Peacock, W.J., (1984) Molecular analysis of the alcohol dehydrogenase (*Adh1*) gene of maize. *Nucleic Acids Research* **12**: 3983-4000.
- Ellis, J.G., Llewellyn, D.J., Dennis, E.S. and Peacock, W.J., (1987) Maize *Adh-1* promoter sequences control anaerobic regulation: addition of upstream promoter elements from constitutive genes is necessary for expression in tobacco. *EMBO Journal* **6**: 11-16.
- Ellis, J.G., Llewellyn, D.J., Walker, J.C., Dennis, E.S. and Peacock, W.J., (1987) The ocs element: a 16 base pair palindrome essential for activity of the octopine synthase enhancer. *EMBO Journal* **6**: 3203-3208.
- Gerlach, W.L., Llewellyn, D.J. and Haseloff, J., (1987) Construction of a plant disease resistance gene from the satellite RNA of tobacco ringspot virus. *Nature (London)* **328**: 802-805 (Reviewed pp. 758-759).
- Llewellyn, D.J., Finnegan, E.J., Ellis, J.G., Dennis, E.S. and Peacock, W.J., (1987) Structure and expression of an alcohol dehydrogenase 1 gene from *Pisum sativum* (cv. Greenfeast). *Journal of Molecular Biology* **195**: 115-123.
- Walker, J.C., Llewellyn, D.J., Mitchell, L.E. and Dennis, E.S., (1987) Anaerobically regulated gene expression; molecular adaptations of plants for survival under flooded conditions. *Oxford Surveys of Plant Molecular and Cell Biology* **4**: 71-93.
- Higgins, T.J.V., Newbigin, E.J., Spencer, D., Llewellyn, D.J. and Craig, S., (1988) The sequence of a pea vicilin gene and its expression in transgenic tobacco plants. *Plant Molecular Biology* **11**: 683-695.
- Landsmann, J., Llewellyn, D., Dennis, E.S. and Peacock, W.J., (1988) Organ regulated expression of the *Parasponia andersonii* haemoglobin gene in transgenic tobacco plants. *Molecular and General Genetics* **214**: 68-73.
- Bouchez, D., Tokuhisa, J.G., Llewellyn, D.J., Dennis, E.S. and Ellis, J.G., (1989) The ocs-element is a component of the promoters of several T-DNA and plant viral genes. *EMBO Journal* **8**: 4197-4204.
- Lyon, B.R., Llewellyn, D.J., Huppatz, J.L., Dennis, E.S. and Peacock, W.J., (1989) Expression of a bacterial gene in transgenic tobacco plants confers resistance to the herbicide 2,4-dichlorophenoxyacetic acid. *Plant Molecular Biology* **13**: 533-540.

- Bogusz, D., Llewellyn, D.J., Craig, S., Dennis, E.S., Appleby, C.A. and Peacock, W.J., (1990) Nonlegume hemoglobin genes retain organ-specific expression in heterologous transgenic plants. *Plant Cell* 2: 633-641.
- Singh, K., Dennis, E.S., Ellis, J.G., Llewellyn, D.J., Tokuhisa, J.G., Wahleithner, J.A. and Peacock, W.J., (1990) OCSBF-1, a maize ocs enhancer binding factor: isolation and expression during development. *Plant Cell* 2: 891-903
- Cousins, Y.L., Lyon, B.R. and Llewellyn, D.J., (1991) Transformation of an Australian cotton cultivar: prospects for cotton improvement through genetic engineering. *Australian Journal of Plant Physiology* 18: 481-494.
- Mooney, P.A., Goodwin, P.B., Dennis, E.S. and Llewellyn, D.J., (1991) *Agrobacterium tumefaciens*-gene transfer into wheat tissues. *Plant Cell Tissue and Organ Culture* 25: 209-218.
- Ellis, J.G., Tokuhisa, J.G., Llewellyn, D.J., Bouchez, D., Singh, K., Dennis, E.S. and Peacock, W.J., (1993) Does the ocs-element occur as a functional component of the promoters of plant genes? *Plant Journal* 4: 433-443.
- Foley, R.C., Grossman, C., Ellis, J.G., Llewellyn, D.J., Dennis, E.S., Peacock, W.J. and Singh, K.B., (1993) Isolation of a maize bZIP protein subfamily: candidates for the ocs element transcription factor. *Plant Journal* 3: 669-679.
- Lyon, B.R., Cousins, Y.L., Llewellyn, D.J. and Dennis, E.S., (1993) Cotton plants transformed with a bacterial degradation gene are protected from accidental spray drift damage by the herbicide 2,4-dichlorophenoxyacetic acid. *Transgenic Research* 2: 162-169.
- Chen, Z., Llewellyn, D.J., Fan, Y., Li, S., Guo, S., Jiao, G. and Zhao, J., (1994) 2,4-D resistant transgenic cotton plants produced by *Agrobacterium* - mediated gene transfer. *Scientia Agricultura Sinica* 27: 31-37.
- Fitt, G.P., Mares, C.L. and Llewellyn, D.J., (1994) Field evaluation and potential ecological impact of transgenic cottons (*Gossypium hirsutum*) in Australia. *Biocontrol Science and Technology* 4: 535-548.
- Llewellyn, D., Cousins, Y., Mathews, A., Hartweck, L. and Lyon, B., (1994) Expression of *Bacillus thuringiensis* insecticidal protein genes in transgenic crop plants. *Agriculture Ecosystems and Environment* 49: 85-93.
- Grover A. , Llewellyn D. and Dennis E.S., (1995) Modification of vacuolar chitinase and osmotin cDNAs of tobacco for extracellular secretion and their cloning with CaMV 35S promoter. *Journal of Plant Biochemistry and Biotechnology* 4:41-42.
- Andersson, C.R., Jensen, E.O., Llewellyn, D.J., Dennis, E.S. and Peacock, W.J., (1996) A new hemoglobin gene from soybean: a role for hemoglobin in all plants. *Proceedings of the National Academy of Science, USA* 93: 5682-5687.
- Harcourt, R.L., Llewellyn, D., Morton, R., Dennis, E.S. and Peacock, W.J., (1996) Effectiveness of purified *Bacillus thuringiensis* Berliner insecticidal proteins in controlling three insect pests of Australian eucalypt plantations. *Journal of Economic Entomology* 89: 1392-1398.

- Llewellyn, D. and Fitt, G., (1996) Pollen dispersal from two field trials of transgenic cotton in the Namoi Valley, Australia. *Molecular Breeding* **2**: 157-166.
- Mathews, A., Llewellyn, D.J., Wu, Y. and Dennis, E.S., (1996) Isolation and characterisation of full length cDNA clones of the giant taro (*Alocasia macrorrhiza*) trypsin/chymotrypsin inhibitor. *Plant Molecular Biology* **30**: 1035-1039.
- Andersson, C.R., Llewellyn, D.J., Peacock, W.J. and Dennis, E.S. (1997) Cell-specific expression of the promoters of two nonlegume hemoglobin genes in a transgenic legume, *Lotus corniculatus*. *Plant Physiology* **113**: 45-57.
- Hartweck, L.M., Llewellyn, D.J. and Dennis, E.S., (1997) The *Arabidopsis thaliana* genome has multiple divergent forms of phosphoinositol-specific phospholipase C. *Gene* **202**: 151-15633.
- Last, D.I. and Llewellyn, D.J., (1997) Antifungal proteins from seeds of Australian native plants and isolation of an antifungal peptide from *Atriplex nummularia*. *New Zealand Journal of Botany* **35**:385-394.
- Hughes, P., Llewellyn, D. and Whitecross, M. (1997) Isolation of a beta-purothionin cDNA from hexaploid wheat. *Plant Physiology* (Electronic Plant Gene Register) **114**: 156.
- Mullins, K.V., Llewellyn, D.J., Hartney, V.J., Strauss, S. and Dennis, E.S. (1997) Regeneration and transformation of *Eucalyptus camaldulensis*. *Plant Cell Reports* **16**: 787-791.
- Murray, F.R., Llewellyn, D.J., Peacock, W.J. and Dennis, E.S. (1997) Isolation of the glucose oxidase gene from *Talaromyces flavus* and characterisation of its role in the biocontrol of *Verticillium dahliae*. *Current Genetics* **32**: 367-375.
- Trevaskis, B., Watts, R.A., Andersson, C.R., Llewellyn, D.J., Hargrove, M.S., Olson, J.S., Dennis, E.S. and Peacock, W.J. (1997) Two hemoglobin genes in *Arabidopsis thaliana*: the evolutionary origins of leghemoglobins. *Proceedings of the National Academy of Science, USA* **94**: 12230-12234.
- Wu, Y., Llewellyn, D., Mathews, A. and Dennis, E.S. (1997) Adaptation of *Helicoverpa armigera* (Lepidoptera: Noctuidae) to a proteinase inhibitor expressed in transgenic tobacco. *Molecular Breeding* **3**: 371-380.
- Kazan, K., Murray, F.R., Goulter, K.C., Llewellyn, D.J. and Manners, J.M. (1998) Induction of cell death in transgenic plants expressing a fungal glucose oxidase. *Molecular Plant Microbe Interactions* **11**: 555-562.
- Southerton, S.G., Strauss, S.H., Olive, M.R., Harcourt, R.L., Decroocq, V., Zhu, X., Llewellyn, D.J., Peacock, W.J. and Dennis, E.S. (1998) Eucalyptus has a functional equivalent of the *Arabidopsis* floral meristem identity gene LEAFY. *Plant Molecular Biology* **37**: 897-910.
- Decroocq, V., Zhu, X., Kauffman, M., Kyojuka, J., Peacock, W.J., Dennis, E.S. and Llewellyn, D.J. (1999) A TM3-like MADS-box gene from *Eucalyptus* expressed in both vegetative and reproductive tissues. *Gene* **228**: 155-160.
- Last, D.I. and Llewellyn, D.J. (1999) Targeted expression of a detoxification gene in transgenic tobacco confers tolerance to 2,4-D. *Weed Science* **47**:401-404.

- Murray, F., Llewellyn, D., McFadden, H., Last, D., Dennis, E.S. and Peacock, W.J. (1999) Expression of the *Talaromyces flavus* glucose oxidase gene in cotton and tobacco reduces fungal infection, but is also phytotoxic. *Molecular Breeding* 5: 219-232.
- Hughes, P., Dennis, E., Whitecross, M., Llewellyn, D. and Gage P. (2000) The cytotoxic plant protein, beta-purothionin, forms ion channels in lipid membranes. *J. Biol. Chem.* 275:823-827.
- Harcourt, R.L. Kyojuka, J. Floyd, R.B. Bateman, K.S. Tanaka, H. Decroocq, V. Llewellyn, D.J. Zhu, X. Peacock W. J. and Dennis E.S. (2000) Insect- and herbicide-resistant transgenic eucalypts. *Molecular Breeding* 6: 307-315.
- Llewellyn, D. (2000) Herbicide tolerant forest trees. In "Molecular Biology of Woody Plants" Vol. 2. S.M. Jain & S.C. Minocha (eds) Kluwer Academic Publishers. Dordrecht, The Netherlands, pp439-466.
- Ellis, M, Millar, A, Llewellyn, DJ, Peacock, WJ, and Dennis ES. (2000) Transgenic cotton (*Gossypium hirsutum*) over-expressing alcohol dehydrogenase show increased ethanol fermentation but no increase in tolerance to oxygen deficiency. *Australian J. Plant Physiology* 27: 1041-1050.
- Ruan, Y-L., Furbank, R.T. and Llewellyn, D.J. (2000) Pathway and control of sucrose import into initiating fibre cells. *Australian J. Plant Physiol.* 27: 795-800.
- Ruan, Y-L. Llewellyn, D.J. and Furbank, R.T. (2001) The control of single-celled cotton fiber elongation by developmentally reversible gating of plasmodesmata and coordinated expression of sucrose and K⁺ transporters and expansin. *Plant Cell* 13: 47-60.
- Gubler, F., Chandler, P., White, R., Llewellyn, D. and Jacobsen, J. (2002) Gibberellin signaling in barley aleurone cells. Control of SLN1 and GAMYB expression. *Plant Physiology* 129: 191-200.
- Llewellyn, D.J and Higgins, T.J.V (2002) Transgenic crop plants with increased tolerance to insect pests. In *Plant Biotechnology and Transgenic Plants* (K-M. Oksman-Caldentey and W. Barz (eds)) Marcel Dekker, New York, pp571-596.
- Rungis, D., Llewellyn, D., Dennis, E.S. and B.R. Lyon, B.R. (2002) Investigation of the chromosomal location of the bacterial blight resistance gene present in an Australian cotton (*Gossypium hirsutum* L.) cultivar. *Australian J. Agricultural Research* 53: 551-560.
- Townsend, B.J., and Llewellyn, D.J. (2002) Spatial and temporal regulation of a soybean (*Glycine max*) lectin promoter in transgenic cotton (*Gossypium hirsutum*). *Functional Plant Biology* 29: 835-843.
- Wu, Y., Llewellyn, D.J., and Dennis, E.S. (2002) A quick and easy method for isolating good quality RNA from cotton (*Gossypium hirsutum* L.) *Plant Molecular Biology Reporter* 20: 213-218.
- Ruan, Y-L., Llewellyn, D.J. and Furbank, R.T. (2003) Suppression of sucrose synthase gene expression represses cotton fiber cell initiation, elongation and seed development. *Plant Cell* 15: 952-964.
- Schünmann, PHD, Llewellyn, D.J., Surin, B., Boevink, P., De Feyter, R.C. and Waterhouse, P.M. (2003) A suite of novel promoters and terminators for plant biotechnology. *Functional Plant Biology* 30: 443-452.

- Rungis, D., Llewellyn, D., Dennis, E.S. and Lyon, B.R. (2005) Simple sequence repeat (SSR) markers reveal low levels of polymorphism between cotton (*Gossypium hirsutum* L.) cultivars of Australian and American origin. *Australian Journal of Agricultural Research* **56**: 301-307.
- Ruan, Y.-L., Llewellyn, D.J., Furbank, R.T. and Chourey, P.S. (2005) The delayed initiation and slow elongation of fuzz-like short fibre cells in relation to altered patterns of sucrose synthase expression and plasmodesmata gating in a lintless mutant of cotton. *Journal of Experimental Botany* **56**: 977-984.
- Townsend B.J., Poole A., Blake C.J., and Llewellyn D.J. (2005) Antisense suppression of a (+)- δ -cadinene synthase gene in cotton prevents the induction of this defense response gene during bacterial blight infection but not its constitutive expression. *Plant Physiology* **138**: 516-528.
- Wu, Y., Rozenfeld, S., Defferrard, A., Ruggiero, K., Udall, J.A., Kim, H.-R., Llewellyn, D.J. and Dennis, E.S. (2005) Cycloheximide treatment of cotton ovules alters the abundance of specific classes of mRNAs and provides a method of generating novel ESTs for microarray expression profiling. *Molecular Genetics and Genomics* **274**: 477-493.
- Wu, Y., Machado, A., White, R.G., Llewellyn, D.J. and Dennis, E.S. (2006) Identification of early genes expressed during cotton fibre initiation using cDNA microarrays. *Plant Cell Physiology* **47**: 107-127.
- Udall, J.A., Swanson, J., Haller, K., Rapp, R.A., Sparks, M.E., Hatfield, J., Yu, Y., Wu, Y., Arpat, A.B., Sickler, B.A., Wilkins T.A., Guo, J.-Y., Chen, X.-Y., Scheffler, J., Taliencio, E., Turley, R., McFadden, H., Payton, P., Allen, R., Zhang, D., Haigler, C., Wilkerson, C., Suo, J., Schulze, S.R., Pierce, M.L., Essenberg, M., Kim, H., Llewellyn, D.J., Dennis, E.S., Kudrna, D., Wing, R., Paterson, A.H., Soderlund, C., and Wendel J.F (2006) A global assembly of cotton ESTs. *Genome Research* **16**: 441-450.
- Amarasinghe, B.H.R.R., Faivre Nitschke, E., Wu, Y., Udall, J.A., Dennis, E.S., Constable, G. and Llewellyn, D.J., (2006) Genomic approaches to the discovery of promoters for sustained expression in cotton (*Gossypium hirsutum* L.) under field conditions: expression analysis in transgenic cotton and Arabidopsis of a Rubisco small subunit promoter identified using EST sequence analysis and cDNA microarrays. *Plant Biotechnology* **23**: 437-450.
- Wu Y., Llewellyn D.J., White R., Ruggiero K., Al-Ghazi Y. and Dennis E.S., (2007). Laser capture microdissection and cDNA microarrays used to generate gene expression profiles of the rapidly expanding fibre initial cells on the surface of cotton ovules. *Planta* **226**: 1475-1490.
- Townsend B.J. and Llewellyn D.J. (2007). Reduced terpene levels in cottonseed add food to fiber. *TRENDS in Biotechnology* **25**: 239-241.
- Lovell D., Wu Y., White R., Machado A., Llewellyn D.J., Dennis E.S. and Anderssen R.S., (2007). Phenotyping cotton ovule fibre initiation with spatial statistics. *Australian Journal of Botany* **55**: 608-617.
- Llewellyn D.J., Mares C.L. and Fitt G.P., (2007). Field performance and seasonal changes in the efficacy against *Helicoverpa armigera* (Hübner) of transgenic cotton expressing the insecticidal protein vip3A. *Agricultural and Forest Entomology* **9**: 93-101.

Llewellyn D., Tyson C., Constable G., Duggan B., Beale S. and Steel P. (2007). Containment of regulated genetically modified cotton in the field. *Agriculture Ecosystems and Environment* **121**: 419-429.

Horvath D.P., Llewellyn D. and Clay S.A., (2007). Heterologous hybridization of cotton microarrays with Velvetleaf (*Abutilon theophrasti*) reveals physiological responses due to corn competition. *Weed Science* **55**: 546-557.

Charles G.W., Constable G.A., Llewellyn D.J. and Hickman M.A. (2007). Tolerance of cotton expressing 2,4-D detoxification gene to 2,4-D applied in the field. *Australian Journal of Agricultural Research* **58**: 780-787.

Ruan Y-L., Llewellyn D.J., Liu Q., Xu S-M., Wu L-M., Wang L. and Furbank, R.T., (2008). Expression of sucrose synthase in the developing endosperm is essential for early seed development in cotton. *Functional Plant Biology* **35**: 382-393.

Dennis E.S., Ellis J., Green A., Llewellyn D., Morell M., Tabe L. and Peacock W.J. (2008). Genetic contributions to agricultural sustainability. *Philosophical Transactions of the Royal Society B* **363**: 591-609.

Machado A., Wu Y., Yang Y., Llewellyn D. and Dennis E.S., (2009). The MYB transcription factor GhMYB25 regulates early fibre and trichome development. *The Plant Journal* **59**: 52-62.

Lacape J-M., Jacobs J., Arioli T., Derijcker R., Forestier-Chiron N., Llewellyn D., Jean J., Thomas E. and Viot C., (2009). A new interspecific, *Gossypium hirsutum* x *G. barbadense*, RIL population: towards a unified consensus linkage map of tetraploid cotton. *Theoretical and Applied Genetics* **119**: 281-292.

Christianson J.A., Wilson I.W., Llewellyn D.J. and Dennis E.S., (2009). The low oxygen-induced NAC domain transcription factor ANAC102 affects viability of Arabidopsis seeds following low-oxygen treatment. *Plant Physiology* **149**: 1724-1738.

Al-Ghazi Y., Bourot S., Arioli T., Dennis E.S. and Llewellyn D.J., (2009). Transcript profiling during fibre development identifies pathways in secondary metabolism and cell wall structure that may contribute to cotton fibre quality. *Plant and Cell Physiology* **50**: 1364-1381.

Christianson J.A., Llewellyn D.J., Dennis E.S. and Wilson I.W., (2010). Global gene expression responses to waterlogging in roots and leaves of cotton (*Gossypium hirsutum* L.). *Plant and Cell Physiology* **51**: 21-37.

Christianson J.A., Dennis E.S., Llewellyn D.J. and Wilson I.W., (2010). ATAF NAC transcription factors: Regulators of plant stress signaling. *Plant Signaling and Behavior* **5**: 428-432.

Paterson A.H., Rong J-K., Gingle A.R., Chee P.W., Dennis E.S., Llewellyn D., Dure L.S. III, Haigler C., Myers G.O., Peterson D.G., ur Rahman M., Zafar Y., Reddy U., Saranga Y., Stewart J.M., Udall J.A., Waghmare V.N., Wendel J.F., Wilkins T.A., Wright R.J., Zaki W., Hafez E.E. and Zhu J. (2010) Sequencing and utilization of the *Gossypium* genomes. *Tropical Plant Biology* **3**: 71-72.

Christianson J.A., Llewellyn D.J., Dennis E.S. and Wilson I.W., (2010). Comparisons of early transcriptome responses to low-oxygen environments in three dicotyledonous plant species. *Plant Signaling and Behavior* **5**: Issue 8, epub ahead of print.

Hinz M., Wilson I.W., Yang J., Buerstenbinder K., Llewellyn D., Dennis E.S., Sauter M., and Dolferus R., (2010) Arabidopsis *RAP2.2*: An ethylene response transcription factor that is important for hypoxia survival. *Plant Physiol.* **153**: 757-772.

Lacape J-M., Llewellyn D., Jacobs J., Arioli T., Becker D., Calhoun S., Al-Ghazi Y., Liu S., Palai O., Georges S., Giband M., de Assunção H., Barroso P.A.V., Claverie M., Gawryziak G., Jane J., Vialle M. and Viot C., (2010) Meta-analysis of cotton fiber quality QTLs across diverse environments in a *Gossypium hirsutum* x *G. barbadense* recombinant inbred line population. *BMC Plant Biology* **10**: 132.

Granted Patents and Patent Applications:

WO2005095614-A1; AU2005229157-A1; US2008196120-A1 Altering fiber initiation and/or elongation in a fiber producing plant comprises manipulating the plant by modifying a transcription factor, regulatory protein, or a cell cycle protein, produced at or around anthesis.

WO200245485-A; WO200245485-A1; US2002116736-A1; AU200220363-A; EP1349446-A1; BR200115782-A; CN1479569-A; ZA200303930-A; AU2002220363-B2; CN1326996-C; US7498492-B2; IN200300565-P2 Altering fibre development or properties of a fibre producing plant by modulating sucrose synthase activity and/or expression in such plants, useful for enhancing fibre yield and quality and for increasing seed size.

US5837849-A Isolated DNA that activates or enhances transcription from plant genes - effective in mono- or di-cotyledonous plants, particularly for increasing expression of heterologous genes.

US5710267-A DNA construct containing plant enhancer bound by octopine synthase transcription factor - and a heterologous plant promoter and gene, useful for generating transgenic plants for increased production of heterologous proteins.

AU9539013-A; AU715462-B Eucalyptus reproductive genes - useful for prodn. of sterile Eucalyptus trees useful for establishing wood lot plantations or in re-forestation projects.

WO9521924-A; EP746616-A; WO9521924-A1; AU9516611-A; EP746616-A1; EP746616-A4; AU692228-B; US6054318-A New glucose oxidase gene constructs - used for transforming organisms, esp. plants, to increase resistance to pests or diseases.

EP278659-A2; EP278659-A; ZA8800319-A; JP63276492-A; CA1309365-C; EP278659-B1; DE3853840-G; ES2074435-T3 Transcription activating element for plant genes - obtd. from octopine synthase (OCS) gene of T-DNA from *Agrobacterium tumefaciens*.

Listing of Claims:

1-116. (Cancelled)

117. A method of increasing fibre initiation and/or elongation in a fibre producing plant comprising genetically manipulating the plant such that the production of a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38 is increased when compared to a wild-type fiber producing plant.
118. A method of reducing fibre initiation and/or elongation in a fibre producing plant comprising genetically manipulating the plant such that the production of a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38 is reduced when compared to a wild-type fiber producing plant.
119. The method of claim 117 or 118, wherein the polypeptide comprises consecutive amino acids whose sequence is set forth in SEQ ID NO: 12.
120. The method of claim 117, wherein the genetic manipulation comprises exposing the plant to a vector which comprises a nucleotide sequence encoding a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38.
121. The method of claim 118, wherein the genetic manipulation comprises exposing the plant to an antisense

Applicant: Yingru Wu, et al.
Serial No.:10/594,785
Filed: September 17, 2007
Exhibit B

polynucleotide or a catalytic polynucleotide which hybridizes to an mRNA molecule encoding the polypeptide, and/or exposing the plant to a dsRNA molecule that specifically down-regulates levels of an mRNA molecule encoding the polypeptide, such that the level of the polypeptide produced by the plant is reduced.

122. The method of claim 121, wherein the genetic manipulation is exposing the plant to the dsRNA molecule and the dsRNA molecule is double-stranded over at least 19 basepairs whose sequence corresponds to a consecutive sequence set forth in SEQ ID NO: 38, or to a consecutive sequence which is identical to the sequence set forth in SEQ ID NO: 38.
123. The method of claim 117 or 118, wherein the plant is a species of the Genus *Gossypium*.
124. A process of assessing the potential of a fibre producing plant to produce fibre, the process comprising analyzing whether the plant has a polynucleotide comprising a sequence identical to the nucleotide sequence set forth in SEQ ID NO: 38, and/or analyzing whether the plant is capable of expressing a polypeptide comprising an amino acid sequence which is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38.
125. The method of claim 124, wherein the polypeptide comprises consecutive amino acids whose sequence is set forth in SEQ ID NO: 12.
126. A substantially purified and/or recombinant polypeptide selected from the group consisting of a polypeptide

comprising consecutive amino acids whose sequence is set forth in SEQ ID NO: 12.

127. An isolated and/or exogenous polynucleotide comprising a polynucleotide selected from the group consisting of:

- i) a polynucleotide comprising consecutive nucleotides whose nucleotide sequence is set forth in SEQ ID NO: 38; and
- ii) a polynucleotide which encodes a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38.

128. A polynucleotide which is:

- i) a catalytic polynucleotide capable of cleaving a polynucleotide whose sequence is identical to the nucleotide sequence set forth in SEQ ID NO: 38, or
- ii) a dsRNA molecule comprising a polynucleotide which is double-stranded over at least 19 basepairs whose sequence corresponds to a consecutive sequence set forth in SEQ ID NO: 38, or to a consecutive sequence which is identical to the sequence set forth in SEQ ID NO: 38.

129. A vector comprising or encoding the polynucleotide of claim 127.

130. A vector comprising or encoding the polynucleotide of claim 128.
131. A plant or bacterial cell comprising the vector of claim 129.
132. A host cell comprising the vector of claim 130.
133. A transgenic plant, the plant having been transformed with the polynucleotide of claim 127.
134. A transgenic plant, the plant having been transformed with the polynucleotide of claim 128.
135. The transgenic plant of claim 133, which when compared to an isogenic non-transgenic plant, produces a modified level of a polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38.
136. The transgenic plant of claim 135, wherein the polypeptide comprises consecutive amino acids whose amino acid sequence is set forth in SEQ ID NO:12.
137. A transgenic seed of the plant of claim 133.
138. A transgenic seed of the plant of claim 134.
139. A process for producing fibre comprising obtaining the transgenic plant of claim 133 so as to thereby produce the fibre.
140. A process of breeding a fibre producing plant having a

polypeptide comprising consecutive amino acids whose sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38, the process comprising analyzing the plant for a genetic variation in a polynucleotide whose sequence is identical to the nucleotide sequence set forth in SEQ ID NO: 38, and/or analyzing the plant for a genetic variation in a polypeptide which is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38, and breeding the fibre producing plant.

141. A process of selecting from a breeding population a fibre producing plant with altered fibre initiation and/or elongation potential, the method comprising:

- i) crossing two plants which have differing potential to produce fibre so as to produce progeny plants,
- ii) performing on the progeny plants a process comprising analyzing the plant for a genetic variation in a polynucleotide whose sequence is identical to the nucleotide sequence set forth in SEQ ID NO: 38, and/or analyzing the plant for a genetic variation in a polypeptide whose amino acid sequence is identical to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 38, and
- iii) selecting a progeny plant with altered fibre initiation and/or elongation potential when compared to a parent plant.

142. A process for identifying an agent which reduces fibre

initiation and/or elongation of a fibre producing plant,
the method comprising:

- i) exposing a polynucleotide comprising consecutive nucleotides whose sequence is identical to the nucleotide sequence set forth in SEQ ID NO: 38 to a candidate agent, and
- ii) assessing the ability of the candidate agent to hybridize and/or-cleave the polynucleotide.